



Care of Mouth after CEREC Crown

A CEREC crown is a restoration that is completed in one visit to our office thanks to the advanced technology we use.

1. Avoid chewing in the area of the crown or drinking hot beverages until any numbness has completely worn off.
2. Gum irritation or sensitivity can occur with any anesthetic injection and crown prep procedure. You may rinse with warm salt water as needed (1 tsp salt mixed with 1 cup warm water).
3. Mild tooth sensitivity to cold can occur with placement of any bonded dental restoration. This should gradually diminish after a few days.
4. We adjust your bite immediately after placing the crown, however, sometimes the numb feeling does not allow for proper refinement. If this occurs, call our office to schedule a time to come in for an adjustment. The need for adjustments to the bite surfaces is higher for patients who have received their treatment under sedation.
5. We will inform you if we find evidence of advanced tooth decay or cracks during your treatment. These can transmit bacteria to the nerve of the tooth which can cause the nerve to develop an infection or die. If this occurs, you may need Root Canal treatment. Signs and symptoms include: persistent or throbbing pain that increases in intensity over time, severe bite sensitivity, pain that radiates to the side of the face or pain that wakes you during the night. Please call the office if you experience any of these symptoms.
6. To keep your CEREC Crown in its best condition, brush and floss regularly and visit us for your routine hygiene care. The most common reason for failure of any restoration is failure to maintain both proper hygiene and regular preventive visits.

Companion Signature: _____ Date: _____
(If patient uses sedation, companion must sign.)

Dr. Brian Heck, DDS, PA | **Dr. Clay Beshore, DDS** | **Dr. Cody Welding, DDS** | **Dr. Kevin Cattaneo, DDS**

5120 Congressional Circle, Suite 100, Lawrence, KS 66049
Office: (785) 856-8550